

Serial No 091 630,333
Group No 1646



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 468 497 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **91112504.5**

(51) Int. Cl.⁵: **C07K 7/06, A61K 37/02**

(22) Date of filing: **25.07.91**

(30) Priority: **26.07.90 US 558031**

(43) Date of publication of application:
29.01.92 Bulletin 92/05

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **MERRELL DOW
PHARMACEUTICALS INC.
2110 East Galbraith Road
Cincinnati Ohio 45215(US)**

(72) Inventor: **Krstenansky, John L.
3455 Rambow Drive
Palo Alto, California 94306(US)**

(74) Representative: **Vossius & Partner
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)**

(54) **Peptide-antagonists of bombesin/GRP.**

(57) A method for controlling the growth of tumor tissues, especially small cell lung and prostatic carcinomas, and gastric acid secretion, causative and symptomatic of, peptic (esophageal, gastric, and duodenal) ulcers. Treatment comprises administering to a patient in need thereof, an effective amount of a bombesin/GRP type inhibitor.

Antagonists of bombesin/GRP which are derivatives of naturally occurring bombesin/GRP possessing a thiomethylene or methylene sulfoxide bond connecting the two amino acids on the carboxy terminal end is modified are described. The antagonism is confirmed using conventional competitive binding and biochemical assays as well as conventional physiological tests and the use of these derivatives in a variety of conditions in which bombesin/GRP is implicated is also described.

EP 0 468 497 A2

This invention relates to novel antimitotic and antisecretory peptides that are antagonists to the related peptides Bombesin and Gastrin Releasing Peptide which are useful in controlling; (1) the growth of small cell lung and prostatic carcinomas, and (2) gastric acid secretions, causative and symptomatic of peptic (esophageal, gastric, and duodenal) ulcers.

Bombesin is a 14 amino acid peptide, originally isolated from the skin of the frog *Bombina bombina*. Bombesin is known to have a range of effects including induction of gastric and pancreatic secretion, stimulation of the nervous system, induction of myoelectric and contractile activity of intestinal myocytes, reduction of renal blood flow, secretion of pituitary hormones, and growth promotion.

Gastrin Releasing Peptide (GRP), a 27 amino acid peptide, was isolated later in mammals, and found to have an identical carboxy terminus to bombesin, with analogous properties. Further, gastrin releasing peptide and the structurally related bombesin have been shown to compete for binding to high affinity receptors on small cell lung carcinoma cells (SCLC), and prostatic endothelial cells. Binding with any of these growth factors causes measurable [³H]thymidine uptake and clonal growth.

Many studies have implicated cell growth factors in tumor cell growth by activation of growth factor receptors. In this regard, Bombesin/GRP and receptor activation is related to other well-known growth factors and oncogenes. There has been considerable interest in the design and development of competitive bombesin receptor antagonist as possible antimitotic agents since the discovery that these peptides, bombesin and GRP, act as potent autocrine growth factors in human small cell lung and possibly also prostatic carcinoma systems. Underlying support for this approach comes from experiments utilizing anti-bombesin monoclonal antibodies to prevent binding of their receptor and eliciting a mitogenic response. Experiments using monoclonal antipeptide antibodies demonstrated clonal growth inhibition, of SCLC cells *in vitro* and in xenografts of nude mice *in vivo*. The mitogenic response begins with the production of intracellular signals by the ligand-receptor complex. For GRP/bombesin, this includes activation of a G-protein, which in turn activates phospholipase C (PLC). PLC converts phosphatidylinositol phosphate (PI) into inositol 1,4,5-triphosphate (IP₃) and diacylglycerol (DAG). IP₃ and DAG are believed to be intracellular signals of the GRP/bombesin receptor pathway activation.

After Bombesin/GRP cell receptors were established on SCLC cells, receptors were also found to be present on human prostate cells. Subsequent studies also firmly established the growth effects of bombesin/GRP on prostatic epithelial cells *in vitro* utilizing bombesin/GRP antibodies. Further, cell proliferation studies of PC3 (a human prostatic carcinoma cell line) and PMU23 (a epithelial cell line from a primary culture of a stage III prostatic carcinoma) demonstrated a dose dependent Bombesin/GRP growth curve. This data suggest that there may be an autocrine or paracrine enhancement of tumor cell growth within the prostate gland. Therefore, interrupting the effect of GRP/bombesin maybe useful for treating the progression of prostatic cancers where these factors are acting as autocrine or paracrine mitotic agents.

Bombesin and GRP are also capable of inducing gastric and pancreatic secretion and have demonstratable saturable receptors on cells of pancreatic (B-Cells) and intestinal (C-cells) origin, and therefore, antagonist of these receptors may serve in treatment of peptic ulcers and intestinal and pancreatic disorders, and may extend to adenocarcinomas of these tissues.

Studies with the anti-bombesin/GRP antibodies lead to the hypothesis that it may be possible to disrupt the autocrine growth cycle of bombesin/GRP using designed peptide receptor antagonists. Since then several types of Bombesin antagonists have been reported. These antagonists have been defined by type and position of the substitutions of the natural sequence. Early receptor antagonists suffered from low potency, lack of specificity, and toxicity which presented serious problems with their scientific and therapeutic use.

More recent work has concentrated on modification of the carboxy terminal (C-terminal) region of these peptides to interrupt the receptor interaction utilizing a variety of different types of C-terminal modified analogs. These have included incorporation of D-amino acids, non-peptide bonds for example (Ψ[CH₂NH]), amide, and ester modifications. These alterations gave rise to certain peptides having improved characteristics.

The applicants have prepared peptide derivatives having N-terminal alkanoyl groups of from 6 to 18 carbon atoms which have markedly improved antagonistic properties relative to the bombesin receptor. The peptide antagonists of this invention potentially possess significant antimitotic and/or antisecretory activity and therefore may allow for a scientifically interesting and therapeutically significant adjunct to cancer therapy and the treatment of ulcers.

Claimed are peptide derivatives of the formula

X-A₁-A₂-A₃-A₄-A₅-A₆-A₇-Y

wherein

- X is a straight, branched, or cyclic alkanoyl group of from 6 to 18 carbon atoms;
- A₁ is Gln, His, his, MeHis, Mehis, His(rMe), his(rMe), His(ΠMe), his(ΠMe), Ala, Val, Leu, leu, MeLeu, Meleu, Ile, Pro, Phe, phe, Met, Trp, Lys, Arg, aNpa, Glu(OMe), glu(OMe), pGlu, pglu, Glu, or glu;
- 5 A₂ is Trp, MeTrp, Ser, Thr, Asn, Gln, Tyr, Met, Phe(Sub), Lys, Lys(pClCbz) or Cha,;
- A₃ is Ala, MeAla, Aib, Phe, phe, Phe(Sub), phe(Sub), bNpa, Arg, Glu, Gln, Gly, Asn, Cys, Ser, Thr, Tyr, Val, Leu, Ile, Pro, Met, Trp, Nle, or Abu;
- A₄ is Val, MeVal, Gln, Gly, Asn, Cys, Ser, Ala, Leu, Ile, Pro, Phe, Phe(pSub), Met, Trp, Nle, Abu, bNpa, Thr, or Thr(Bzl);
- 10 A₅ is Gly, Ser, ser, Ser(Bzl), ser(Bzl), Thr, Asn, Gln, Tyr, Trp, Phe, phe, Phe(Sub), phe(Sub), Sar, ala, pro, lys, Asp, arg, lys(pClCbz), Ac³c, Ac⁵c, or Ac⁶c;
- A₆ is His, Ser, Thr, Asn, Gln, Tyr, Trp, His, Lys, Lys(Cbz), Arg, Tyr, Phe, phe, Phe(pSub), phe(pSub), His, MeHis, His(rMe), His(ΠMe), Aib, Nlc, Nva, Ser, Ser(Bzl), Glu, Asp, Asp(OtBu), or bNpa;
- A₇ is a bond or is Leu, leu, MeLeu, Meleu, Phe, Ala, Val, MeVal, Ile, Melle, Pro, Met, Trp, Ser, Thr,
- 15 Asn, Gln, Tyr, Trp, Nle, MeNle, Nva, MeNva, or Glu;
- Y is a carboxy terminal residue selected from OH, (C₁-C₈) alkoxy ester, carboxamide, mono or di (C₁-C₄) alkyl amide, (C₁-C₄) alkylamine, (C₁-C₄) thioalkylether,

or a pharmaceutically acceptable salt thereof and certain pharmaceutical uses thereof.

The following common abbreviations of (1) amino acids and their three letter codes, (2) modified and
 20 unusual amino acids, and (3) terminal amino and carboxy substituents used throughout this specification:

(1): THE AMINO ACIDS AND THEIR THREE LETTER CODE

L-AMINO ACIDS

25	Ala	- alanine
	Arg	- arginine
	Asn	- asparagine
	Cys	- cysteine
30	Gly	- glycine
	Glu	- glutamic acid
	Val	- valine
	Gln	- glutamine
	His	- histidine
35	Ile	- isoleucine
	Leu	- leucine
	Lys	- lysine
	Phe	- phenylalanine
	Met	- methionine
40	Pro	- proline
	Ser	- serine
	Thr	- threonine
	Trp	- tryptophan
	Tyr	- tyrosine

(2): MODIFIED AND UNUSUAL AMINO ACIDS

	Ac3c	- 1-amino-1-cyclopropanecarboxylic acid
	Ac5c	- 1-amino-1-cyclopentanecarboxylic acid
50	Ac6c	- 1-amino-1-cyclohexanecarboxylic acid
	Asp(OtBu)	- O-t-butylaspartic acid
	Abu	- α-amino-n-butyric acid
	Aib	- 2-aminoisobutyric acid
	Cha	- cyclohexylalanine
55	Chg	- cyclohexylglycine
	His(ΠMe)	- N ^Π -methyl histidine
	His(rMe)	- N ^r -methyl histidine
	Hyp	- hydroxyproline

	pGlu	- pyroglutamic acid
	glu(OMe)	- gamma-methyl-D-glutamic acid
	Tyr(I)	- 3-iodotyrosine (Tyr(I)),
		3,5-diiodotyrosine (Tyr(I ₂))
5	Lys(Cbz)	- benzyloxycarbonyl lysine
	Lys(pClCbz)	- p-chlorobenzyloxy lysine
	MeLys	- N-methyllysine
	MeLeu	- N-methylleucine
	MeHis	- N-methylhistidine
10	MeTrp	- N-methyltryptophan
	MeAla	- N-methylalanine
	MeVal	- N-methylvaline
	MeNle	- N-methyl(2S)-2-aminohexanoic acid
	MeNva	- N-methyl-(2S)-2-aminopentanoic acid
15	Melle	- N-methylisoleucine
	MePhe	- N-methylphenylalanine
	MePgl	- N-phenylglycine
	aNpa	- α -(naphthyl)alanine
	bNpa	- β -(naphthyl)alanine
20	Mnl	- 3,4-dihydroproline
	Nle	- norleucine
	Nva	- norvaline
	Orn	- ornithine
	Pip	- pipercolate
25	Pba	- p-aminophenylbutyric acid
	Phe(pSub)	- para substituted phenylalanine,
		Phe(pCl) - parachlorophenylalanine,
		Phe(pBr) - parabromophenylalanine,
		Phe(pFl) - parafluorophenylalanine,
30		Phe(pNO ₂) - paranitrophenylalanine
	Pgl	- phenylglycine
	Ser(Bzl)	- benzylserine
	Sar	- sarcosine (N-methylglycine)
	Phe(Sub)	- ortho, meta, or para, mono- or disubstituted phenylalanine
35	Tha	- β -(2-thienyl)-alanine
	Thr(Bzl)	- benzylthreonine
	Trp(Bzl)	- benzyltryptophan
	Tiq	- Tetrahydroisoquinoline 3-carboxylate

40 AMINO AND CARBOXY TERMINAL ACID SUSTITUENTS

	Ac	- acetyl
	Azt	- azetidine-2-carboxylate
	Cin	- cinnamoyl
45	DhCin	- 3,4-dihydrocinnamoyl
	Glt	- glutaryl
	Mal	- maleyl
	Oac	- 8-aminooctanoic acid
	Oct	- n-octyl
50	Suc	- succinyl
	Glt	- glutaryl
	Tfa	- trifluoroacetyl
	#	- C-terminal amide

55 Amino Acids & Modifications

The naturally occurring amino acids, with the exception of glycine, contain a chiral carbon atom. Unless otherwise specifically indicated, the optically active amino acids, referred to herein, are of the L-configuration.

tion. For example, any of the amino acids of the A₁ or A₇ group can be of the D- or L-configuration. According to convention the code for a particular amino acid beginning with a lower case letter indicates the amino acid is of the D-configuration, while a code beginning with an upper case letter can indicate either the L-configuration or both the D-and L-configuration as context indicates. As is customary, the structure of peptides written out herein is such that the amino terminal end is on the left side of the chain and the carboxy terminal end is on the right side of the chain.

An alkanoyl group of from 6 to 18 carbon atoms is taken to include straight, branched, cyclic, saturated and unsaturated acyl groups optionally having from 1 to 4 double bonds, for example, n-hexanoyl, isohexanoyl, cyclohexanoyl, cyclopentylcarbonyl, heptanoyl, n-octanoyl, n-decanoyl, n-undecanoyl, n-dodecanoyl, n-tridecanoyl, n-tetra-decanoyl, 3,7-dimethyloctanoyl, 2,4-dimethylhexadecanoyl, 1,7,10-trimethylundecanoyl, n-pentadecanoyl, n-hexadecanoyl, eicosoyl, n-heptadecanoyl, 3-propylnonanoyl, 10-undecanoyl, 3,7,11-trimethyl-2,6,10-pentadecatrienoyl, 5,9-dimethyl-2,4,8-decatrienoyl, 4,6-dimethyloctanoyl, 1,2,5,9-tetramethyl-2,4,8-decatrienoyl, 2-hexadecanoyl, lauroyl, myristicoyl, palmitoyl, oleicoyl, linoleicoyl, γ -linolenicoyl, and stearoyl.

The term "any amino acid" as used herein does not purport to include any carboxylic acid having an amino substituent, but rather is used as it is commonly used by those skilled in the art of polypeptide derivatives and includes the naturally occurring amino acids as well as other "non-protein" α -amino acids commonly utilized by those in the peptide chemistry arts when preparing synthetic analogs of naturally occurring peptides. The naturally occurring amino acids are glycine, alanine, valine, leucine, isoleucine, serine, methionine, threonine, phenylalanine, tyrosine, tryptophan, cysteine, proline, histidine, aspartic acid, asparagine, glutamic acid, glutamine, arginine, ornithine, and lysine. Also included would be the D-isomers of the naturally occurring amino acids; D-alanine, D-valine, D-leucine, D-isoleucine, D-serine, D-methionine, D-threonine, D-phenylalanine, D-tyrosine, D-tryptophan, D-cysteine, D-proline, D-histidine, D-aspartic acid, D-asparagine, D-glutamic acid, D-glutamine, D-arginine. Also included are "non-protein" α -amino acids, examples are 1-amino-1-cyclopropanecarboxylic acid (Ac3c), 1-amino-1-cyclopentanecarboxylic acid (Ac5c), 1-amino-1-cyclohexanecarboxylic acid (Ac6c), O-t-butylaspartic acid (Asp(OtBu)), α -amino-n-butyric acid (Aba), (2S)-2-aminohexanoic acid (norleucine)(Nle), (2S)-2-aminopentanoic acid (norvaline)(Nva), N¹-methylhistidine (His(IIMe)), N⁷-methylhistidine (His(τ Me)), N-methyllysine (MeLys), N-methylleucine (MeLeu), N-methylhistidine (MeHis), N-methyltryptophan (MeTrp), N-methylalanine (MeAla), N-methylvaline (MeVal), N-methyl-(2S)-2-aminohexanoic acid (MeNle), N-methyl-(2S)-2-aminopentanoic acid (MeNva), N-methylisoleucine (Melle), N-methylphenylalanine (MePhe), N-methylphenylglycine (MePgl), β -(2-thienyl)-alanine (Tha), benzylthreonine (Thr(Bzl)), benzyltryptophan (Trp(Bzl)) norleucine, norvaline, alloisoleucine, homoarginine, thiaproline, dihydropyridine, hydroxyproline (Hyp), homoserine, cyclohexylglycine (Chg), α -amino-n-butyric acid (Aba), cyclohexylalanine (Cha), aminophenylbutyric acid (Pba), phenylalanines mono or disubstituted at the ortho, meta, or para, such as para substituted phenylalanine (Phe(pSub)) and para-chlorophenylalanine, para-aminophenylalanine and para-nitrophenylalanine or positions of the phenyl moiety with one or two of the following, a (C₁-C₄) alkyl, (C₁-C₄) alkoxy, halogen, or nitro groups or substituted with a methylenedioxy group, β -2- and 3-thienylal-alanine, β -2- and 3-furanylalanine, β -2-, 3-, and 4-pyridylalanine, β -(benzothienyl-2- and 3-yl)alanine, β -(1-and 2-naphthyl)alanine(Npa), O-alkylated derivatives of serine, threonine, or tyrosine, methyl esters of glutamic and aspartic acid, S-alkylated cysteine, the O-sulfate ester of tyrosine, and halogenated tyrosines such as 3-iodotyrosine, and 3,5-diiodotyrosine.

Groups of α -amino acids can be defined by certain charge characteristics. There are two general characteristics of side chains: nonpolar and polar. The nonpolar residues are made up of the hydrophobic residues which includes those with aliphatic hydrocarbon side chains: Gly, Ala, Val, Leu, Ile, Nle, Pro and the aromatic group Phe and Trp, and the pseudohydrocarbon, Met. The polar amino acids are made up of three groups: (1) The acidic hydrophobic residues, (2) the neutral residues, and (3) basic hydrophobic residues. The neutral residues contain the hydroxyl-containing residues, Ser and Thr; the amides, Asn and Gln; aromatic rings, Tyr and Trp; the sulfhydryl groups, Cys, and small structurally accommodating amino acids Ala and Gly. In the polar class are "acidic hydrophilic" residues that include aspartic acid, glutamic acid, and tyrosine and the "basic hydrophilic" residues that include His, Lys, and Arg.

Y designates the chemical group(s) that may be utilized to substitute or modify the terminal amino acid. Therefore, Y may be a carboxamide, alkoxy ester, alkylamide, alkyl amine, or thioalkylether.

The polypeptides of formula 1 can form pharmaceutically acceptable salts with any non-toxic, organic or inorganic acid. Illustrative inorganic acids which form suitable salts include hydrochloric, hydrobromic, sulphuric and phosphoric acid and acid metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Illustrative organic acids which form suitable salts include the mono, di and tricarboxylic acids. Illustrative of such acids are, for example, acetic, glycolic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic, benzoic, hydroxybenzoic,

phenylacetic, cinnamic, salicylic, 2-phenoxybenzoic and sulfonic acids such as methane sulfonic acid and 2-hydroxyethane sulfonic acid. Salts of the carboxy terminal amino acid moiety include the non-toxic carboxylic acid salts formed with any suitable inorganic or organic bases. Illustratively, these salts include those of alkali metals, as for example, sodium and potassium; alkaline earth metals, such as calcium and magnesium; light metals of Group IIIA including aluminum; and organic primary, secondary and tertiary amines, as for example, trialkylamines, including triethylamine, procaine, dibenzylamine, 1-ethenamine, N,N'-dibenzylethylenediamine, dihydroabietylamine, N-(lower)alkylpiperidine, and any other suitable amine.

The ability of the peptide derivatives of this invention to act as antagonists of Bombesin/GRP can be demonstrated by the ability of such peptides to compete with radioiodinated bombesin/GRP for mammalian bombesin/GRP receptors using the method of Buck, et al., *Science* 226: 987-989, 1984, and by the ability of such compounds to stimulate or to inhibit bombesin/GRP induced phosphatidylinositol turnover using the method of Bristow, et al., *British J. Pharmacol.* 90: 211-21, 1987.

As with any generic group of chemical compounds, certain groups are preferred. Applicants prefer those peptide derivatives of formula 1 wherein:

- X is a straight, branched, or cyclic alkanoyl group of from 6 to 18 carbon atoms;
- A₁ is Gln, His, his, MeHis, Mehis, His(τ Me), his(τ Me), His(TMe), or his(TMe);
- A₂ is Trp or MeTrp;
- A₃ is Ala MeAla, Val, Leu, or Ile;
- A₄ is Val, MeVal, Gly, Ala, Leu, Ile, or Nle;
- A₅ is Gly or ala;
- A₆ is His; MeHis, His(τ Me), or His(TMe);
- A₇ is a bond or is Leu, leu, MeLeu, Meleu, Ala, Val, MeVal, Ile, MeLie, Nle, MeNle, Nva, or MeNva; and
- Y is a carboxy terminal residue selected from OH, (C₁-C₈) alkoxy ester, carboxamide, mono or di (C₁-C₄) alkylamide, (C₁-C₄) alkyl amine, (C₁-C₄) thioalkylether, or pharmaceutically acceptable salt thereof.

More preferred are peptide derivatives of formula 1 wherein

- X is an alkanoyl group of from 6 - 10 carbon atoms;
- A₁ is Gln or His;
- A₂ is Trp;
- A₃ is Ala;
- A₄ is Val;
- A₅ is Gly;
- A₆ is His;
- A₇ is a bond or is Leu; and
- Y is a carboxy terminal residue selected from OH or amino.

Most preferred are those compounds of formula 1 wherein

- X is a straight, branched, or cyclic alkanoyl group of from 7 - 9 carbon atoms, especially octanoyl;
- A₁ is Gln or His;
- A₂ is Trp;
- A₃ is Ala;
- A₄ is Val;
- A₅ is Gly;
- A₆ is His;
- A₇ is a bond or is Leu; and
- Y is a carboxy terminal residue selected from OH or amino.

Peptide Synthesis:

The peptides of this invention can be prepared by a variety of procedures readily known to those skilled in the art. Such procedures include the solid phase sequential and block synthesis, gene cloning and combinations of these techniques. The solid phase sequential procedure can be performed using established automated methods such as by use of an automated peptide synthesizer. In this procedure, the peptides were synthesized on the resin beginning with the C-terminal amino acid attached to a methylbenz-hydrylamine resin. The extension of the peptide sequence was done using standard methodology and that of the manufacturer and that known by people skilled in the art.

After completion of coupling of the sequence either the Boc protecting group was left in place or it was removed and the N-terminal amino group acylated. Displacement of the protected fragment from the resin

was accomplished using the hydrogen fluoride procedure.

The α -amino protecting group employed with each amino acid introduced into the polypeptide sequence may be any such protecting group known to the art. Among the classes of α -amino protecting groups contemplated are (1) acyl type protecting groups such as: formyl, trifluoroacetyl, phthalyl, toluenesulfonyl (tosyl), benzenesulfonyl, nitrophenylsulfonyl, tritylsulfonyl, o-nitrophenoxycarbonyl and α -chlorobutyryl; (2) aromatic urethan type protecting groups such as benzyloxycarbonyl and substituted benzyloxycarbonyl, such as p-chlorobenzyloxycarbonyl, p-nitrobenzyl carbonyl, p-bromobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α , α -dimethyl-3,5-dimethoxybenzyloxycarbonyl and benzhydryloxycarbonyl; (3) aliphatic urethan protecting groups such as tert-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl and allyloxycarbonyl; (4) cycloalkyl urethane type protecting groups such as cyclopentylloxycarbonyl, adamantylloxycarbonyl and cyclohexylloxycarbonyl; (5) thiourethane type protecting groups such as phenylthiocarbonyl; (6) alkyl type protecting groups such as triphenylmethyl (trityl) and benzyl; and (7) trialkylsilane groups such as trimethylsilane. The preferred α -amino protecting group is tert-butyloxycarbonyl (Boc).

As is known in the art of solid phase peptide synthesis many of the amino acids bear functionalities requiring protection during the chain preparation. The use and selection of the appropriate protecting group is within the ability of those skilled in the art and will depend upon the amino acid to be protected and the presence of other protected amino acid residues on the peptide. The selection of such a side chain protecting group is critical in that it must be one which is not removed by cleavage during cleavage of the protecting group of the α -amino moiety. For example, suitable side chain protecting groups for lysine are benzyloxycarbonyl and substituted benzyloxycarbonyl, said substituent being selected from halo (e.g., chloro, bromo, fluoro) and nitro (e.g., 2-chlorobenzyloxycarbonyl, p-nitrobenzyloxy-carbonyl, 3,4-dichlorobenzyloxycarbonyl), tosyl, t-amylloxycarbonyl, t-butyloxycarbonyl and diisopropylmethoxycarbonyl. The alcoholic hydroxyl group of threonine and serine can be protected with an acetyl, benzoyl, tert-butyl, trityl, benzyl, 2,6-dichlorobenzyl or benzyloxycarbonyl group. The preferred protecting group is benzyl.

The selection of an appropriate coupling reagent is within the skill of the art. A particularly suitable coupling reagent where the amino acid to be added is Gln, Asn or Arg is N,N'-diisopropylcarbodiimide and 1-hydroxybenzotriazole. The use of these reagents prevents nitrile and lactam formation. Other coupling agents are (1) carbodiimides (e.g., N,N'-dicyclohexylcarbodiimide and N-ethyl-N'-(γ -dimethylaminopropyl)carbodiimide); (2) cyanamides (e.g., N,N-dibenzylcyanamide); (3) ketenimines; (4) isoxazolium salts (e.g., N-ethyl-5-phenyl-isoxazolium-3'-sulfonate); (5) monocyclic nitrogen containing heterocyclic amides of aromatic character containing one through four nitrogens in the ring such as imidazolidines, pyrazolidines, and 1,2,4-triazolidines. Specific heterocyclic amides that are useful include N,N'-carbonyldiimidazole and N,N-carbonyl-di-1,2,4-triazole; (6) alkoxylated acetylene (e.g., ethoxyacetylene); (7) reagents which form a mixed anhydride with the carboxyl moiety of the amino acid (e.g., ethylchloroformate and isobutylchloroformate) or the symmetrical anhydride of the amino acid to be coupled (e.g., Boc-Ala-O-Ala-Boc), (8) nitrogen containing heterocyclic compounds having a hydroxy group on one ring nitrogen (e.g., N-hydroxyphthalimide, N-hydroxysuccinimide and 1-hydroxybenzotriazole), and (9) diphenyl phosphorylazide. Other activating reagents and their use in peptide coupling are described by Kapoor, J. Pharm. Sci., 59, pp. 1-27 (1970). Applicants prefer the use of the symmetrical anhydride as a coupling reagent for all amino acids except Arg, Asn and Gln.

Each protected amino acid or amino acid sequence is introduced into the solid phase reactor in about a four-fold excess and the coupling is carried out in a medium of dimethylformamide: methylene chloride (1:1) or in dimethylformamide alone or preferably methylene chloride alone. In cases where incomplete coupling occurs, the coupling procedure is repeated before removal of the α -amino protecting group, prior to the coupling of the next amino acid in the solid phase reactor. The success of the coupling reaction at each stage of the synthesis is monitored by the ninhydrin reaction as described by E. Kaiser et al, *Analyt. Biochem.* 34, 595 (1970).

Following the coupling of the α -amino protected amino acid to the resin support, the protecting group is removed using any suitable procedure such as by using trifluoroacetic acid in methylene chloride, trifluoroacetic acid alone, or HCl in dioxane. The deprotection is carried out at a temperature of between 0°C and room temperature. Other standard cleaving reagents and conditions for removal of specific α -amino protecting groups may be used. After removal of the α -amino protecting group the other amino protected amino acids are coupled step-wise in the desired order. Alternatively, multiple amino acid groups may be coupled by the solution method prior to coupling with the resin supported amino acid sequence.

After the desired amino acid sequence has been obtained, the peptide is removed from the resin. This can be done by hydrolysis such as by treatment of the resin bound polypeptide with an amino acid alcohol and acetic acid in dichloromethane (DCM). Protecting groups can be removed by procedures well known in

the art. Typically protecting group removal is done after the peptide chain synthesis is complete but the protecting groups can be removed at any other appropriate time. Purification of peptides is principally accomplished through preparative reversed phase high performance liquid chromatography and those techniques known to those skilled in the art.

Therapeutic Use:

The natural history of peptic ulcer disease is one of recurrent exacerbations and remissions. As a result, ulcerative diseases should be treated as a chronic disorder. Peptic (esophageal, gastric, and duodenal) ulcers occur in areas of the gastrointestinal tract exposed to acid and pepsin. The compounds of the present invention, which are useful in the treatment of gastrointestinal and/or pancreatic ulcers and may be effective in resultant hypersecretions occurring from the pancreas and/or stomach, particularly hydrochloric acid and pepsin. As such compounds of this invention may serve as an appropriate intervention to treat peptic ulcers.

The appropriate dose of a peptide derivative of this invention when used in the treatment of peptic ulcers is from 0.2 mg/kg to 250 mg/kg of patient body weight per day depending on factors such as the patient, the severity of the peptic ulcer to be treated and the peptide derivative selected. The suitable dose for a particular patient can be readily determined. Preferably from 1 to 4 daily doses would be administered typically with from 5 mg to 100 mg of active compound per dose. The amount of a peptide of this invention required to inhibit gastric acid secretion can be readily determined by those skilled in the art.

Use of bombesin/GRP analogs in cancer therapy is indicated for the treatment of SCLC and prostatic carcinomas and prevention of a variety of other cancer conditions. Those experienced in this field are readily aware of the circumstances requiring cancer therapy. The term "patient" used herein is taken to mean mammals such as primates, including humans, sheep, horses, cattle, pigs, dogs, cats, rats and mice.

As used herein, the term "tumor tissue" means both benign and malignant tumors or neoplasms and includes melanomas, lymphomas, leukemias, and sarcomas. Illustrative examples of tumor tissues are cutaneous such as malignant melanomas and mycosis fungoides; hematologic tumors such as leukemias, for example, acute lymphoblastic, acute myelocytic, or chronic myelocytic leukemia; lymphomas such as Hodgkin's disease or malignant lymphoma; gynecologic tumors such as ovarian and uterine tumors; urologic tumors such as those of the prostate, bladder, or testis; soft tissue sarcomas, osseous, or non-osseous sarcomas, breast tumors; tumors of the pituitary, thyroid, and adrenal cortex; gastrointestinal tumors such as those of the esophagus, stomach, intestine, and colon; pancreatic and hepatic tumors; laryngeal papillomatoses and lung tumors.

The term "controlling the growth" and the concept of treating a cancer means slowing, interrupting, arresting, or stopping the growth and metastases of a rapidly proliferating tumor in a warm blooded animal; it being understood that treatment in a warm blooded animal does not generally provide a "cure" for the tumor in the sense that necessarily the tumor tissue is destroyed or totally eliminated.

The appropriate dose of a peptide derivative of this invention when used in the treatment of, or controlling the growth of a cancer or tumor tissue, including small cell lung carcinoma or prostatic cancers is from 0.2 mg/kg to 250 mg/kg of patient body weight per day depending on factors such as the patient, the severity of the small cell lung carcinoma to be treated and the peptide derivatives selected. The suitable dose for a particular patient can be readily determined. Preferably from 1 to 4 daily doses would be administered typically with from 5 mg to 100 mg of active compound per dose. The amount of a peptide of this invention required to inhibit SCLC growth can be readily determined by those skilled in the art.

It is generally known that therapeutic agents used in the treatment of cancer can be used in conjunction with other therapeutic agents or therapies known to be useful in the treatment of cancer. The peptides of invention can also be used in conjunctive therapy. For example, a peptide of structure 1 can be administered in conjunction with surgical excision of the tumor or with radiation therapy, immunotherapy, or local heat therapy. In the preferred mode of treating cancer a peptide of structure 1 are administered in conjunction with a chemical cytotoxic agent known to be useful for tumor therapy. Illustrative chemical cytotoxic agents are cyclophosphamide, methotrexate, prednisone, 6-mercaptopurine, procarbazine, danorubicin, chlorambucil, cytosine arabinoside, 6-thioguanine, thio TEPA, 5-fluorouracil, 5-fluoro-2-deoxyuridine, 5-azacytidine, nitrogen mustard, 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU), 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea (CCNU), busulfan, adriamycin, bleomycin, vindesine, cycloheximide, or methylglyoxal bis(guanylhydrazine) (MGBG). When such combination therapy is employed, the effect of the cytotoxic agent may be potentiated. The remission produced by the cytotoxic agent may be enhanced and regrowth of the tumor may be slowed or prevented. Use of such combination therapy therefore allows for smaller doses or fewer individual doses of the cytotoxic agent to be employed. The term "combination therapy"

contemplates the administration of a peptide of structure 1 immediately prior to the beginning of therapy with a cytotoxic agent, concurrently with such therapy, or during the time immediately following the cessation of such therapy. When such combination therapy is employed for the treatment of a tumor the cytotoxic agent may be administered at a dosage known in the art to be effective for treating the tumor.

5 However, a peptide of structure 1 may produce an additive or synergistic effect with a cytotoxic agent against a particular tumor. Thus when such combination antitumor therapy is used, the dosage of the cytotoxic agent administered may be less than that administered when the cytotoxic agent is used alone. In combination with a peptide derivative of structure 1, the cytotoxic agent may, therefore, be administered at a lower dosage or at less frequent intervals as compared to the cytotoxic agent used alone. Similarly when
10 combination therapy is used, the dosage or frequency of administration of a peptide of structure 1 may be less than when used without a cytotoxic agent.

Although some of the peptide derivatives may survive passage through the gut following oral administration, applicants prefer non-oral administration, for example, subcutaneous, intravenous, intramuscular or intraperitoneal; administration by depot injection; by implant preparation; or by application to the mucous
15 membranes, such as, that of the nose, throat and bronchial tubes, for example, in an aerosol can containing a peptide derivative of this invention in a spray or dry powder form.

For parenteral administration the compounds may be administered as injectable dosages of a solution or suspension of the compound in a physiologically acceptable diluent with a pharmaceutical carrier which can be a sterile liquid such as water and oils with or without the addition of a surfactant and other
20 pharmaceutically acceptable adjuvants. Illustrative of oils which can be employed in these preparations are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, and mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions, ethanol and glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions.

25 EXAMPLES

This invention is illustrated by the following, nonlimiting examples.

EXAMPLE 1

30 PREPARATION OF

N-(n-Octanoyl)-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

The peptide was synthesized on p-methylbenzhydrylamine resin using N^t-t-Boc protected amino acids
35 and octanoic acid that were coupled with the Bop reagent (ref. Dung Le-Nguyen, Annie Heitz and Bernard Castro, J. Chem. Soc., Perkin Trans. I (1987) 1915-1919. Upon completion of the synthesis, the peptide was deprotected and removed from the resin by treatment with liquid HF containing 10% anisole for 1 hour at 0° C. Upon removal of the HF *in vacuo*, the peptide was precipitated with Et₂O and filtered. The peptide was then extracted from the resin with 30% aqueous acetic acid and glacial acetic acid. The wash was
40 lyophilized to give the crude peptide, which was purified by HPLC. The peptide was analyzed by amino acid analysis, fast-atom bombardment-mass spectroscopy and analytical HPLC.

In a similar manner the compounds of Examples 2 - 4 are prepared.

EXAMPLE 2

45 N-(Lauroyl)-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

EXAMPLE 3

50 N-(Palmitoyl)-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

EXAMPLE 4

N-(n-Octanoyl)-Gln-Trp-Ala-Val-ala-His-NH₂

55 The compounds of examples 1 - 4 have the following physical and chemical properties.

Compound (Example #)	Amino Acid Analysis					FAB-MS (M + H) ⁺
	Glx	Ala	Val	Leu	His	
1	1.01(1)	2.01(2)	0.97(1)	1.02(1)	0.99(1)	949.5
2	1.03(1)	2.00(2)	0.96(1)	1.02(1)	0.98(1)	1005.5
3	1.04(1)	1.98(2)	0.98(1)	1.03(1)	0.98(1)	1061.5
4	1.09(1)	1.94(2)	0.88(1)	--	1.09(1)	836.4

EXAMPLE 5

AFFINITY FOR THE BOMBESIN RECEPTOR AS DEMONSTRATED BY COMPETITION AT THE RECEPTOR FOR IODINATED GRP

The pancreata from one or more mice were pooled, minced, and homogenized in 50 mM HEPES (pH 7.4) containing 120 mM NaCl and 5 mM KCl at 4°C and centrifuged at 37,500 X g for 15 minutes. The pellet was resuspended in 50 mM HEPES (pH 7.4) containing 10 mM EDTA and 300 mM KCl and incubated for 30 minutes at 4°C. The suspension was centrifuged as above and the pellet was washed two times in 50 mM HEPES (pH 7.4) containing protease inhibitors (10 uM phenylmethylsulfonyl fluoride, 1 uM thiorphan, 40 ug/ml bacitracin, 10mM iodoacetic acid and 4 ug/ml leupeptin), and again centrifuged. The tissue was then resuspended in incubation buffer (1ml per 4 mg pancreas) and incubated for 15 minutes at room temperature, then 250 ul was added to each assay tube to commence the assay. The assay tubes contained incubation buffer consisting of 50 mM HEPES (pH 7.4), 0.5% BSA, protease inhibitors, 2 mM MnCl₂, 1 uM somatostatin, and concentrations of ¹²⁵I-GRP and peptides as needed in a final volume of 500 ul. The assay was allowed to proceed to equilibrium for 90 minutes at room temperature. After this time, the contents of each tube was rapidly filtered over Whatman GF/B filters presoaked in 0.1% polyethyleneimine and the filters were rapidly washed three times with ice-cold 50 mM HEPES (pH 7.4). Filter-bound radioactivity was quantitated in a gamma counter. Competition of iodinated GRP binding by test compounds or standards was expressed as a percentage of ¹²⁵I-GRP binding in the absence of peptide. Affinity and maximal binding were calculated with Ligand (Biosoft, Cambridge, UK).

EXAMPLE 6

ANTAGONISM OF THE BOMBESIN RECEPTOR AS DEMONSTRATED BY THE EFFECT ON PHOSPHATIDYLINOSITOL TURNOVER

Pancreata from mice were pooled and chopped at 350 μm with a tissue chopper. The chopped tissue was washed twice with Krebs-Hepes, then incubated for 30 minutes in 37°C Krebs-Hepes buffer with fresh buffer after 15 minutes. The tissue was then incubated in this buffer containing 100-200 μCi of ³H-inositol at 37°C for 1 hour. The tissue was then washed twice and incubated for another 30 minutes in Krebs-Hepes (containing 10 mM Li⁺) at 37°C with fresh buffer change after 15 minutes. Portions of the tissue mass (approximately 10 mg per assay tube) were then placed in Li⁺ buffer with protease inhibitors, (40 ug/ml bacitracin, 4 ug/ml leupeptin, 4 ug/ml chymostatin, 20 ug/ml thiorphan), 0.1% BSA, and then 0.1-1000 nM peptide was added in 25 μl in a final volume of 250 μl. To measure antagonism portions of the tissue in Li⁺ buffer were pretreated with 1 uM of the potential antagonist for 5 minutes at 25°C before the addition of agonist (GRP). After 60 minutes at room temperature, the phosphatidylinositol turnover was terminated by the addition of 940 μl chloroform:methanol (1:2), followed by 310 μl chloroform, followed by 310 μl water. Each tube was then vortexed for 5 seconds and then centrifuged at 2500 x g for 8 minutes to separate the phases. 900 μl of the top (aqueous) phase was then mixed with 2.1 ml water and loaded onto a 0.5 ml Biorad AG-1X8 (formate) ion exchange column. 50 μl of the bottom phase (chloroform) was withdrawn from each tube and placed in a counting vial, dried, and counted in scintillation fluid. The material on the columns was washed in order with: 1) 10 ml of water 2) 5 ml of 5 mM disodium tetraborate/60 mM sodium

formate 3) 10 ml of 1 M ammonium formate in 0.1 M formic acid. The final (third) wash was collected and one ml was mixed with 10 ml of Bio-Safe scintillant and counted. The ratio of these counts (total inositol phosphates) to the corresponding organic phase counts was then calculated for each sample. The ratios in the presence of test compound and/or standards were then compared to the ratios for control tubes (i.e., no stimulating agonist). Dose-response lines were constructed and the abilities of test compounds to stimulate or to inhibit GRP induced phosphatidylinositol turnover were determined by graphical analysis or with the aid of a computer program.

Following the procedures of examples 5 and 6 with the compounds of examples 1 - 4, the following data was obtained.


Compound (Example #)	Bombesin Receptor Binding ¹ (Example 5)	Antagonism
1	++	+
2	+	+
3	+	+
4	++	+

¹ + indicates $K_D \leq 500 \text{ nM}$

++ indicates $K_D \leq 50 \text{ nM}$

Claims

1. A peptide derivative of the formula



wherein

- X is a straight, branched, or cyclic alkanoyl group of from 6 to 18 carbon atoms;
- A₁ is Gln, or His;
- A₂ is Trp;
- A₃ is Ala;
- A₄ is Val;
- A₅ is Gly, or ala;
- A₆ is His;
- A₇ is a bond, or is Leu;
- Y is a carboxy terminal residue selected from OH, (C₁-C₈) alkoxy ester, carboxamide, mono or di (C₁-C₄) alkyl amide, (C₁-C₄) alkylamine, (C₁-C₄) thioalkylether, or pharmaceutically acceptable salt thereof.

2. A peptide derivative of claim 1 wherein

- X is a straight, branched, or cyclic alkanoyl group of from 6 to 10 carbon atoms;
- A₁ is Gln or His;
- A₂ is Trp;
- A₃ is Ala;
- A₄ is Val;
- A₅ is Gly;
- A₆ is His;
- A₇ is a bond or is Leu; and
- Y is a carboxy terminal residue selected from OH or amino.

3. A peptide derivative of claim 1 wherein

- X is a straight, branched, or cyclic alkanoyl group of from 7 - 9 carbon atoms;
- A₁ is Gln or His;
- A₂ is Trp;
- A₃ is Ala;

A₄ is Val;
 A₅ is Gly;
 A₆ is His;
 A₇ is a bond or is Leu; and
 5 Y is a carboxy terminal residue selected from OH or amino.

4. A peptide derivative of claim 1 wherein

X is octanoyl;
 A₁ is Gln or His;
 10 A₂ is Trp;
 A₃ is Ala;
 A₄ is Val;
 A₅ is Gly;
 A₆ is His;
 15 A₇ is a bond or is Leu; and
 Y is a carboxy terminal residue selected from OH or amino.

5. A peptide derivative of claim 1 which is

X-Gln-Trp-Ala-Val-ala-His-NH₂

20 wherein X is octanoyl.

6. A peptide derivative of claim 1 which is

X-Gln-Trp-Ala-Val-ala-His-NH₂

25 wherein X is lauroyl or palmitoyl.

7. A peptide derivative of claim 1 which is

X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

wherein X is octanoyl.

8. A peptide derivative of claim 1 which is

X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

30 wherein X is lauroyl.

9. A peptide derivative of claim 1 which is

35 X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂

wherein X is palmitoyl.

10. A process for preparing a peptide derivative according to any one of claims 1 to 9 comprising the steps of:

- 40 a) Using a resin with a suitably bound C-terminal protected amino acid from the group A₇;
 b) sequentially coupling the other alpha amino protected amino acids, A₆ through A₁, to achieve the protected amino acid sequence claimed;
 c) removing said protecting groups and purifying the desired peptide.

11. Pharmaceutical composition containing a compound according to any one of claims 1 to 9 optionally in combination with a pharmaceutically acceptable carrier and/or diluent.

12. Pharmaceutical composition according to claim 11 for the treatment of small cell lung carcinoma or conditions associated with small cell lung carcinoma.

13. Pharmaceutical composition according to claim 11 for the treatment of prostatic carcinoma or conditions associated with prostatic carcinoma.

14. Pharmaceutical composition according to claim 11 for controlling the growth of tumor tissue or for the treatment of conditions associated with the growth of tumor tissue.

15. Pharmaceutical composition according to claim 11 for the treatment of peptic ulcers or conditions associated with peptic ulcers.


16. Pharmaceutical composition according to claim 11 for effecting antagonism of bombesin/GRP or for the treatment of conditions associated with effecting antagonism of bombesin/GRP.

17. Pharmaceutical composition according to claim 11 for the treatment of cancer or conditions associated with cancer.

18. Pharmaceutical composition according to claim 11 for the treatment of conditions associated with platelet aggregation.

10 **Claims for the following Contracting State: ES**

1. A process for preparing a peptide derivative of the formula



15 wherein

X is a straight, branched, or cyclic alkanoyl group of from 6 to 18 carbon atoms;

A₁ is Gln, or His;

A₂ is Trp;

20 A₃ is Ala;

A₄ is Val;

A₅ is Gly, or ala;

A₆ is His;

A₇ is a bond or is Leu;

25 Y is a carboxy terminal residue selected from OH, (C₁-C₈) alkoxy ester, carboxamide, mono or di (C₁-C₄) alkyl amide, (C₁-C₄) alkylamine, (C₁-C₄) thioalkylether,

or a pharmaceutically acceptable salt thereof comprising the steps of:

a) using a resin with a suitably bound C-terminal protected amino acid from the group A₇;

30 b) sequentially coupling the other alpha amino protected amino acids, A₆ through A₁, to achieve the protected amino acid sequence claimed;

c) removing said protecting groups and purifying the desired peptide.

2. A process of preparing a peptide derivative according to claim 1 wherein

35 X is a straight, branched, or cyclic alkanoyl group of from 6 to 10 carbon atoms;

A₁ is Gln or His;

A₂ is Trp;

A₃ is Ala;

A₄ is Val;

A₅ is Gly;

40 A₆ is His;

A₇ is a bond or is Leu; and

Y is a carboxy terminal residue selected from OH or amino.

3. A process of preparing a peptide derivative according to claim 1 wherein

45 X is a straight, branched, or cyclic alkanoyl group of from 7 - 9 carbon atoms;

A₁ is Gln or His;

A₂ is Trp;

A₃ is Ala;

A₄ is Val;

50 A₅ is Gly;

A₆ is His;

A₇ is a bond or is Leu; and

Y is a carboxy terminal residue selected from OH or amino.

55 4. A process of preparing a peptide derivative according to claim 1 wherein

X is octanoyl;

A₁ is Gln or His;

A₂ is Trp;

A₃ is Ala;
 A₄ is Val;
 A₅ is Gly;
 A₆ is His;
 5 A₇ is a bond or is Leu; and
 Y is a carboxy terminal residue selected from OH or amino.

5. A process of preparing a peptide derivative according to claim 1 wherein the product obtained is
 X-Gln-Trp-Ala-Val-ala-His-NH₂
 10 wherein X is octanoyl.
6. A process of preparing a peptide derivative according to claim 1 wherein the product obtained is
 X-Gln-Trp-Ala-Val-ala-His-NH₂
 wherein X is lauroyl or palmitoyl.
 15
7. A process of preparing a peptide derivative according to claim 1 wherein the product obtained is
 X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂
 wherein X is octanoyl.
- 20 8. A process of preparing a peptide derivative according to claim 1 wherein the product obtained is
 X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂
 wherein X is lauroyl
9. A process of preparing a peptide derivative according to claim 1 wherein the product obtained is
 25 X-Gln-Trp-Ala-Val-ala-His-Leu-NH₂
 wherein X is palmitoyl.
10. A process for the preparation of a pharmaceutical composition which comprises combining a com-
 pound obtained according to any one of claims 1 to 9 with a pharmaceutically acceptable carrier and/or
 30 diluent.
11. A process according to claim 10 wherein the pharmaceutical composition obtained is used for the
 treatment of small cell lung carcinoma or conditions associated with small cell lung carcinoma.
- 35 12. A process according to claim 10 wherein the pharmaceutical composition obtained is used for the
 treatment of prostatic carcinoma or conditions associated with prostatic carcinoma.
13. A process according to claim 10 wherein the pharmaceutical composition obtained is used for
 controlling the growth of tumor tissue or for the treatment of conditions associated with the growth of
 40 tumor tissue.
14. A process according to claim 10 wherein the pharmaceutical composition obtained is used for the
 treatment of peptic ulcers or conditions associated with peptic ulcers.
- 45 15. A process according to claim 10 wherein the pharmaceutical composition obtained is used for effecting
 antagonism of bombesin/GRP or for the treatment of conditions associated with effecting antagonism of
 bombesin/GRP.
16. A process according to claim 10 wherein the pharmaceutical composition obtained is used for the
 50 treatment of cancer or conditions associated with cancer.
17. A process according to claim 10 wherein the pharmaceutical composition obtained is used for the
 treatment of conditions associated with platelet aggregation.

55



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 468 497 A3**

EUROPEAN PATENT APPLICATION

Application number: **91112504.5**

Int. Cl.⁵: **C07K 7/06, A61K 37/02**

Date of filing: **25.07.91**

Priority: **26.07.90 US 558031**

Date of publication of application:
29.01.92 Bulletin 92/05

Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

Date of deferred publication of the search report:
06.05.92 Bulletin 92/19

Applicant: **MERRELL DOW
PHARMACEUTICALS INC.
2110 East Galbraith Road
Cincinnati Ohio 45215(US)**

Inventor: **Krstenansky, John L.
3455 Rambow Drive
Palo Alto, California 94306(US)**

Representative: **Vossius & Partner
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)**

Peptide-antagonists of bombesin/GRP.

A method for controlling the growth of tumor tissues, especially small cell lung and prostatic carcinomas, and gastric acid secretion, causative and symptomatic of, peptic (esophageal, gastric, and duodenal) ulcers. Treatment comprises administering to a patient in need thereof, an effective amount of a bombesin/GRP type inhibitor.

Antagonists of bombesin/GRP which are derivatives of naturally occurring bombesin/GRP possessing a thiomethylene or methylene sulfoxide bond connecting the two amino acids on the carboxy terminal end is modified are described. The antagonism is confirmed using conventional competitive binding and biochemical assays as well as conventional physiological tests and the use of these derivatives in a variety of conditions in which bombesin/GRP is implicated is also described.

EP 0 468 497 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 11 2504

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	EP-A-0 315 367 (IMPERIAL CHEMICAL INDUSTRIES) 10 May 1989 * page 2, line 45 - page 3, line 46 * * page 14, line 40 - line 54 * * Claim 1, see R1 * ---	1-2, 11-17	C07K7/06 A61K37/02
X	WO-A-9 003 980 (ADMINISTRATORS OF THE TULANE EDUCATIONAL FUND) 19 April 1990 * Claim 19 * ---	1-2, 11-17	
X	EP-A-0 309 297 (THE ADMINISTRATORS OF THE TULANE EDUCATIONAL FUND) 29 March 1989 * page 3, line 18 - page 4, line 5 * ---	1-2, 11-17	
A	US-A-4 331 661 (MARKI ET AL) 25 May 1982 * Abstract * -----	1-18	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C07K A61K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 06 MARCH 1992	Examiner KORSNER S. E.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons ----- Δ: member of the same patent family, corresponding document	